

## Symptom Attenuation by a Satellite RNA *in Vivo* Is Dependent on Reduced Levels of Virus Coat Protein

Jianlong Wang and Anne E. Simon<sup>1</sup>

Department of Biochemistry and Molecular Biology and Program in Molecular and Cellular Biology,  
University of Massachusetts, Amherst, Massachusetts 01003

Received March 19, 1999; returned to author for revision April 14, 1999; accepted April 26, 1999

Many plant RNA viruses provide replication and encapsidation functions for one or more satellite RNAs (sat-RNAs) that can modulate the symptoms of the associated helper virus. Sat-RNA C, a virulent sat-RNA associated with turnip crinkle virus (TCV), normally intensifies symptoms but can attenuate symptoms if the TCV coat protein (CP) is replaced with that of cardamine chlorotic fleck carmovirus [Kong *et al.* (1995) *Plant Cell* 7, 1625–1634] or if TCV contains an alteration in the CP initiation codon (TCV-CP<sub>m</sub>) [Kong *et al.* (1997b) *Plant Cell* 9, 2051–2063]. To further elucidate the mechanism of symptom attenuation by sat-RNA C, the composition of the CP produced by TCV-CP<sub>m</sub> (CP<sub>CP<sub>m</sub></sub>) was determined. Our results reveal that CP<sub>CP<sub>m</sub></sub> likely has two additional amino acids at its N-terminus compared with wild-type TCV CP. TCV-CP<sub>m</sub> produces reduced levels of CP, and this reduction, not the two additional residues at the CP N-terminus, is responsible for symptom attenuation by sat-RNA C. © 1999 Academic Press

**Key Words:** turnip crinkle virus; satellite RNAs; symptom modulation; plant RNA viruses.

### INTRODUCTION


Many plant RNA viruses are associated with one or more nonessential RNAs, such as defective interfering RNAs and satellite RNAs (sat-RNAs). These subviral RNAs depend on the helper virus for replication, encapsidation, and movement through the plant. The sat-RNAs range in size from 194 to 1500 nucleotides (nt) and usually have sequence unrelated to the viral genome. As molecular parasites of their helper viruses, sat-RNAs frequently modulate viral symptom expression. The smaller sat-RNAs (194–700 nt), including those of cucumber mosaic virus (CMV) and turnip crinkle virus (TCV), do not encode any functional open reading frames (ORFs) (Roossinck *et al.*, 1992). However, despite the absence of gene products, these sat-RNAs can have dramatic effects on the symptoms induced by their helper viruses (for reviews, see Kaper and Collmer, 1988; Simon, 1988; Collmer and Howell, 1992). Many viral sat-RNAs attenuate disease (e.g., satellites of CMV and tobacco ringspot virus), a property of interest for viral disease control (Gerlach *et al.*, 1987; Harrison *et al.*, 1987; Tien and Wu, 1991). However, some satellites exacerbate symptoms of their helper virus. One such sat-RNA is sat-RNA C associated with TCV (Li and Simon, 1990).

Some viruses are associated with several sat-RNAs that differentially modulate symptoms (Murant and Kumar, 1990; Blok *et al.*, 1994; Celix *et al.*, 1997). Identical

sat-RNAs can also have different effects on symptoms when associated with different helper virus strains (Kaper *et al.*, 1990; Sleat and Palukaitis, 1990b; Roossinck *et al.*, 1992; Sleat *et al.*, 1994; Kong *et al.*, 1997a; Militao *et al.*, 1998). Furthermore, the same sat-RNA/helper virus combination can have different effects in different hosts. For example, symptoms of CMV are intensified by some CMV sat-RNAs, resulting in either chlorosis in tomato and tobacco (Palukaitis, 1988) or necrosis in tomato (Takanami, 1981; Sleat *et al.*, 1994). Therefore, sat-RNA-mediated symptom modulation is determined by a trilateral interaction among the satellite, helper virus, and host plant. Many sat-RNA sequences that mediate symptom modulation have been defined (Simon *et al.*, 1988; Palukaitis, 1988; Baulcombe *et al.*, 1988; Kurath and Palukaitis, 1989; Masuta and Takanami, 1989; Jaegle *et al.*, 1990; Sleat and Palukaitis, 1990a, 1990b; Naidu *et al.*, 1992; Sleat *et al.*, 1994; Zhang *et al.*, 1994; Oncino *et al.*, 1995; Kong *et al.*, 1997a; Rodriguez-Alvarado and Roossinck, 1997; Taliansky and Robinson, 1997). In contrast, the role of the helper virus and host plant is poorly understood.

We are studying symptom modulation of TCV by sat-RNA C in the host plant *Arabidopsis thaliana*. TCV is a member of the *Carmovirus* genus and is the only genus member that is associated with confirmed sat-RNAs. TCV has a single plus-sense RNA genome of ~4 kb (Carrington *et al.*, 1989; Oh *et al.*, 1995) and two subgenomic (sg) RNAs of 1.72 and 1.45 kb (Carrington *et al.*, 1987; Wang and Simon, 1997) (Fig. 1A). The viral genomic RNA is the mRNA for p28 and its readthrough product p88, which are required for viral rep-

<sup>1</sup>To whom reprint requests should be addressed. Fax: (413) 545-4529. E-mail: [simon@biochem.umass.edu](mailto:simon@biochem.umass.edu).



**FIG. 1.** TCV, TCV-CPm, and associated sgRNAs. (A) Schematic representation of TCV and TCV-CPm genomic RNAs and sgRNAs. The mutation in TCV-CPm is indicated (AUG → ACG). ORFs are denoted by boxes and the encoded products are indicated in the ORFs of the genomic RNA. RdRp, RNA-dependent RNA polymerase; CP, coat protein. ?? indicates the uncertain N-terminus of the TCV-CPm CP. (B) Subviral RNAs associated with TCV. Numbers above the bars indicate analogous positions in the TCV genomic RNA. Sat-RNA C contains nearly full-length sat-RNA D at its 5' end and two regions of TCV at its 3' end.

lication. Two small ORFs (p8 and p9) that are implicated in viral movement (Hacker *et al.*, 1992; Li *et al.*, 1998) can be translated from the 1.72-kb sgRNA *in vitro* (Li *et al.*, 1998). The 1.45-kb sgRNA is the mRNA for the 38-kDa coat protein (CP) (Carrington *et al.*, 1989). The TCV virion is icosahedral in structure with  $T = 3$  symmetry and is composed of 180 copies of the CP (Hogle *et al.*, 1986; Carrington *et al.*, 1989).

TCV systemically infects all ecotypes of *A. thaliana* tested except for ecotype Di-0 (Simon *et al.*, 1992). TCV can replicate in protoplasts of Di-0 but cannot move systemically in whole plants (Simon *et al.*, 1992). When the CP ORF of TCV was replaced with that of the related carmovirus cardamine chlorotic fleck, the resulting chimeric virus (TCV-CP<sub>CCFV</sub>) systemically infected Di-0, indicating that the CP ORF or the CP is an important viral determinant in the resistance of Di-0 to TCV (Oh *et al.*, 1995; Kong *et al.*, 1995). TCV with a single mutation in the initiation codon of the CP ORF (TCV-CPm; Fig. 1A) also overcomes the resistance of Di-0, suggesting that the CP, as opposed to the RNA encoding the CP, is the most likely elicitor of resistance to wild-type (wt) TCV in Di-0.

Sat-RNA C is a hybrid sat-RNA consisting of a nearly full-length avirulent sat-RNA (sat-RNA D) at the 5' end joined to two regions from the 3' end of TCV genomic RNA (Fig. 1B). Sat-RNA C can either intensify or attenuate symptoms depending on the helper virus. Sat-RNA C intensifies symptoms of wt TCV on all hosts where TCV produces visible symptoms (Li and Simon, 1990). In susceptible ecotypes of *A. thaliana*, the moderate stunting symptoms of TCV are exacerbated by sat-RNA C, resulting in the death of the plant by ~16 days postinoculation

(p.i.) (Li and Simon, 1990; Simon *et al.*, 1992). Whole plant *in situ* hybridizations indicate that the TCV genomic RNA becomes concentrated in younger tissue in the presence of sat-RNA C, leading to an inhibition of bolting and plant death (Kong, 1996). On hosts that are tolerant to TCV infection (and therefore have no symptoms associated with virus infection), the presence of sat-RNA C has no effect on plant symptomatology (Li and Simon, 1990).

Although sat-RNA C intensifies the symptoms of wt TCV, it attenuates the symptoms of TCV-CP<sub>CCFV</sub> (Kong *et al.*, 1995). sat-RNA C also attenuates the symptoms of ~70% of plants inoculated with TCV-CPm. Attenuation of symptoms directly correlates with undetectable levels of viral genomic RNA in extracts of whole plants (Kong *et al.*, 1997a, 1997b). Because TCV-CPm differs from TCV by only a single base alteration in the CP initiation codon, the CP and not the RNA encoding the CP is most likely the viral determinant associated with sat-RNA C symptom modulation (Kong *et al.*, 1997a).


Plants and protoplasts infected with TCV-CPm accumulate ~20% of wt levels of a protein that migrates to a similar position as wt TCV CP (CP<sub>WT</sub>) in polyacrylamide gels and cross-reacts with CP-specific antibodies (Kong *et al.*, 1997b). Virions, however, are nearly undetectable in preparations of TCV-CPm-infected protoplasts (Kong *et al.*, 1997b). The inability to isolate comparable levels of virions based on the amount of CP present in cells could be due to CP levels being below the threshold required for virion formation. Alternatively, the TCV-CPm CP (CP<sub>CPm</sub>) may be incapable of efficient virion formation due to translation initiating near, but not at, the natural initiation codon, resulting in a mutant N-terminus. Furthermore, the ability of sat-RNA C to attenuate TCV-CPm symptoms could be due to the presence of reduced levels of CP, the absence of virions, and/or putative alterations to the N-terminus of CP<sub>CPm</sub>.

To distinguish between these possibilities and to further our understanding of symptom modulation by sat-RNA C, the possible initiation codon of CP<sub>CPm</sub> was characterized. Our results suggest that translation of CP<sub>CPm</sub> likely initiates at an in-frame non-AUG codon (CUG) upstream from the normal initiation codon resulting in two additional amino acids at the CP<sub>CPm</sub> N-terminus. In addition, reduced levels of CP, not alterations to the N-terminus, are the principal determinant of symptom attenuation by sat-RNA C. Possible models are proposed to explain how CP levels may be involved in symptom attenuation by sat-RNA C in the TCV/*Arabidopsis* system.

## RESULTS

### TCV-CPm likely initiates translation of its CP from a noncanonical initiation codon resulting in two additional amino acids at its N-terminus

Because CP<sub>WT</sub> and CP<sub>CPm</sub> migrate to similar positions on denaturing polyacrylamide gels, translation of CP<sub>CPm</sub>



**FIG. 2.** Characterization of the initiation codon of TCV-CPm CP. (A) Schematic representation of the CP ORFs in TCV, TCV-CPm (CPm), and the mutants (CPm-L, CPm-T, and CPm-O). The putative initiation codons are boxed, and the mutated nucleotides are indicated by lowercase letters. Two possible CP amino acid sequences are shown for TCV-CPm. The dashed lines in the CPm mutants indicate unaltered sequence. (B) RNA gel blots of total RNA and protein gel blots of total protein and virions accumulating in protoplasts 40 h after inoculation with transcripts of TCV, TCV-CPm (CPm), and CPm mutants as shown in (A). The *A. thaliana* protoplasts ( $5 \times 10^6$ ) were inoculated with  $20 \mu\text{g}$  of wt TCV or mutant transcripts. Total RNA was subjected to RNA gel blot analyses using a probe specific for TCV genomic RNA (gRNA) and sgRNAs (1.72 and 1.45 kb) (see Table 1) or rRNAs (Simon *et al.*, 1992). CP and virions (v) were visualized by chemiluminescence using anti-TCV CP antibody. Each lane represents total protein or virions extracted from  $2.5 \times 10^5$  or  $8.3 \times 10^5$  protoplasts, respectively. Species corresponding to TCV genomic RNA and the 1.45- and 1.72-kb sgRNAs are indicated. (C) Symptoms of Col-0 or Di-0 plants inoculated with TCV, TCV-CPm (CPm), or CPm-L. Seedlings at the six- to eight-leaf stage were inoculated with the wt and mutant transcripts, with (+C) or without sat-RNA C as shown below the plants. Representative plants were photographed at 17 days p.i.. Mock plants were treated with inoculation buffer alone.

was thought to initiate at or near the wt initiation codon. Because the nearest in-frame AUGs in the TCV CP mRNA are either 42 amino acids upstream followed by two stop codons or 40 amino acids downstream from the wt initiation codon and because all nearby out-of-frame AUG triplets are followed by stop codons, CP<sub>CPm</sub> is likely synthesized beginning with a noncanonical initiation codon such as the ACG at the wt initiation position or an in-frame CUG codon located six nucleotides upstream (Fig. 2A). Noncanonical translation initiation has been confirmed or suggested for several cellular mRNAs

(Beams *et al.*, 1991; Saris *et al.*, 1991) and viral RNAs from plant (Schmitz *et al.*, 1996) and animal (Mehdi *et al.*, 1990; Reynolds *et al.*, 1995) hosts (for reviews, see Gallie, 1993; Rohde *et al.*, 1994). Translation was previously reported to initiate from ACG and CUG with 15% and 30% efficiency, respectively, compared with AUG in plant cells (Gordan *et al.*, 1992).

To determine whether the ACG or CUG might serve as initiation codons for CP<sub>CPm</sub>, mutations were introduced into the analogous positions of these nucleotides in TCV-CPm cDNA. As shown in Fig. 2A, the ACG codon

was altered to AGG generating CPm-L and the CUG codon was changed to UUG generating CPm-T. CPm-O was generated by combining both new mutations into a single construct. Neither of the resultant triplets (AGG or UUG) initiates translation in either mammalian or plant cells (Gallie, 1993; Rohde *et al.*, 1994). Transcripts synthesized *in vitro* representing CPm-L, CPm-T, and CPm-O were inoculated onto *A. thaliana* protoplasts, and total RNA, protein, and virions were extracted 40 h p.i. for examination of viral RNA accumulation, CP synthesis, and virion formation.

RNA gel blot analysis revealed that CPm-L and CPm-T replicated to near wt levels, whereas TCV-CPm replicated to a lower level (Fig. 2B) as previously described (Kong *et al.*, 1997b; note, however, that the TCV-CPm sample is underloaded according to the corresponding rRNA levels in Fig. 2B). It is not known why CPm-L and CPm-T had higher levels of genomic RNA replication compared with the parental mutant TCV-CPm. Surprisingly, no viral RNA was detected in CPm-O-infected protoplasts. This was unexpected because the mutations introduced into CPm-O are outside the viral p28 and p88 ORFs and are not within promoters known to be required for replication (Hacker *et al.*, 1992; Wang and Simon, 1997). In addition, previous studies have shown that TCV genomic RNA accumulates in the absence of the majority of the CP ORF, including this region (Kong *et al.*, 1995). However, the possibility cannot be ruled out that there is an additional mutation in some other critical location within the genome of CPm-O.

Western blot analysis using antibodies raised against CP<sub>WT</sub> revealed that protoplasts infected with TCV-CPm accumulated 23% of wt levels of CP and no detectable virions as previously reported (Fig. 2B; Kong *et al.*, 1997b). Protoplasts infected with CPm-L, which maintains the CUG codon but not the ACG codon, accumulated 35% of wt levels of CP. In addition, like TCV-CPm, no virions were detected in CPm-L-infected protoplasts (Fig. 2B). In contrast, no CP was detected in protoplasts infected with CPm-T, which maintains the ACG codon but not the CUG codon (Fig. 2B). As expected, no CP was synthesized in CPm-O-infected protoplasts due to the lack of viral RNA accumulation (Fig. 2B). These results suggest that the CUG is likely the initiation codon for translation of CP<sub>CPm</sub>.

To determine whether CPm-L mimics TCV-CPm in infectivity and symptom modulation by sat-RNA C, *A. thaliana* ecotypes Col-0 and Di-0 were inoculated with TCV, TCV-CPm, and CPm-L, with or without sat-RNA C. Symptoms were assessed visually at various times up to 17 days p.i. As shown in Fig. 2C, CPm-L, like TCV-CPm, was infectious on both Col-0 and Di-0, and unlike TCV, which only infects Col-0. In addition, symptoms of CPm-L were delayed by 1–2 days compared with wt TCV, similar to the symptom delay from inoculation with TCV-CPm. Coinoculation with sat-RNA C resulted in 70% of the plants inoculated with CPm-L remaining either symptomless

(Col-0) or exhibiting attenuated symptoms (Di-0) at 17 days p.i. (Fig. 2C; plants that did not have attenuated symptoms exhibited systemic symptoms similar to those of plants inoculated with only the genomic RNA). No symptoms developed in either CPm-T- or CPm-O-infected plants, with or without sat-RNA C (data not shown). Because TCV requires its CP for systemic movement in plants (Hacker *et al.*, 1992), the inability of CPm-T to infect plants was most likely due to the lack of CP synthesized during the infection.


Taken together, these results suggest that the CUG six nucleotides upstream of the wt initiation codon is the most likely initiation codon used in translating CP<sub>CPm</sub>. Because methionine appears to be the initiating amino acid in all noncanonical translation initiations investigated so far (Gupta and Patwardhan, 1988; Curran and Kolakofsky, 1988; Hann *et al.*, 1988; Peabody, 1989), there probably are two additional amino acids (glutamic acid and threonine) at the N-terminus of CP<sub>CPm</sub> compared with CP<sub>WT</sub> (Fig. 3A). These additional amino acids could account for the slightly slower migration of CP<sub>CPm</sub> compared with CP<sub>WT</sub>, which is visible on most polyacrylamide gels (see Fig. 3B, for example).

#### TCV producing wt levels of CP with two additional N-terminal amino acids abolishes symptom attenuation by sat-RNA C

To determine whether lack of detectable virions and symptom attenuation by sat-RNA C when associated with TCV-CPm is due to the two putative amino acids at the N-terminus of the CPs and/or the reduced levels of CP, the effect of wt levels of mutant CP on symptom modulation and virion formation were investigated. To obtain wt levels of this mutant CP, nucleotides specifying the two extra amino acids likely found in CP<sub>CPm</sub> were inserted after the initiator AUG of the CP ORF in wt TCV generating TCV-CPm3 (Fig. 3A).

Transcripts synthesized *in vitro* were inoculated onto *A. thaliana* protoplasts, and total RNA extracted at 40 h p.i. was subjected to RNA gel blot analysis. As shown in Fig. 3B, TCV-CPm3 replicated to near wt levels (note that the TCV-CPm3 sample was underloaded according to the level of rRNA). Western blot analyses of total protein and virion preparations at 40 h p.i. revealed that TCV-CPm3 produced levels of CP comparable to wt TCV and virions were synthesized (Fig. 3B). The CP of TCV-CPm3 comigrated with CP<sub>CPm</sub> at a position slightly slower than CP<sub>WT</sub> (Fig. 3B), supporting the hypothesis that CP<sub>CPm</sub> contains similar additional amino acids at its N-terminus. Because CP<sub>CPm3</sub> was capable of forming virions, the additional residues at the N-terminus do not by themselves abrogate virion formation.

TCV-CPm3 caused more severe stunting on ecotype Col-0 than wt TCV or TCV-CPm and was able to overcome the resistance of Di-0 to wt TCV (Fig. 3C). Symp-



**FIG. 3.** Effect of increased levels of mutant CP on symptom modulation by sat-RNA C. (A) Schematic representation of the CP ORFs in TCV, TCV-CPm (CPm), and TCV-CPm3 (CPm3). The putative CP initiation codons are in italics. The inserted sequence encoding two additional amino acids in CPm3 are indicated in bold. (B) RNA gel blots of total RNA and protein gel blots of total protein and virions accumulating in protoplasts 40 h after inoculation with transcripts of TCV, TCV-CPm (CPm), and TCV-CPm3 (CPm3). Abbreviations are described in legend to Fig. 2. The experiment was performed as described in Fig. 2. (C) Symptoms of Col-0 and Di-0 plants inoculated with transcripts of TCV, CPm, and CPm3, with (+C) or without sat-RNA C. Inoculation and photography of the plants were performed as described in Fig. 2.

toms developed concomitantly with those of wt TCV (1–2 days earlier than those of TCV-CPm), probably from either increased CP levels or the presence of virions. Although TCV-CPm3 mimicked TCV-CPm in overcoming the resistance of Di-0 to TCV, coinoculation with sat-RNA C neither attenuated nor intensified the symptoms of TCV-CPm3 (Fig. 3C). These results indicate that the N-terminus of the CP is important in the resistance of Di-0 to TCV and suggest that the N-terminal mutations by themselves do not abrogate virion formation or cause symptom attenuation of TCV-CPm by sat-RNA C.

#### Reduced levels of CP<sub>WT</sub> cause symptom attenuation by sat-RNA C

Because the presence of additional amino acids at the N-terminus of the CP do not lead to symptom attenuation by sat-RNA C, the reduced amount of CP synthesized from a noncanonical initiation codon may be the cause of symptom attenuation. If so, then reducing the synthesis

of CP in wt TCV infections should affect sat-RNA C symptom modulation.

To reduce the synthesis of CP<sub>WT</sub>, a GC base pair in the 1.45-kb sgRNA promoter (Wang *et al.*, 1999) was replaced with a weaker GU base pair generating TCV-SGm (Fig. 4A). The mutation did not alter the p9 ORF where the promoter is located. Five percent of wt levels of the 1.45-kb sgRNA was synthesized according to RNA gel blots of total RNA extracted from TCV-SGm-infected protoplasts at 40 h p.i. (Fig. 4B). Western blot analysis of total protein extracted from TCV-SGm-infected protoplasts at 40 h p.i. revealed CP<sub>WT</sub> levels similar to those produced by TCV-CPm (Fig. 4B). This result suggests that translation initiation from the wt initiator AUG is more efficient than the nearby noncanonical initiator CUG. Virions were also markedly reduced in TCV-SGm-infected protoplasts (Fig. 4B), suggesting that the amount of CP accumulating in cells is near a threshold level for virion assembly.

To determine whether TCV-SGm mimics TCV-CPm in



tion, the effect is more pronounced if the CP contains additional amino acids at its N-terminus.

## DISCUSSION

Previous studies (Kong *et al.*, 1995) demonstrated that sat-RNA C was able to attenuate the symptoms of TCV when the CP initiation codon was changed from AUG to ACG (TCV-CPm; Kong *et al.*, 1997b). Because lower levels of possibly mutant CP were produced, it was not clear if the attenuation of symptoms by sat-RNA C was due to the reduced levels of CP or the mutant nature of the CP. To address the nature of CP<sub>CPm</sub>, mutations were introduced into two possible noncanonical in-frame initiation codons in TCV-CPm: the ACG that replaced the wt AUG initiation codon and an upstream in-frame CUG. Converting the ACG to a codon not used for translation initiation (construct CPm-L) did not affect the synthesis of CP, whereas mutating the CUG triplet (construct CPm-T) eliminated CP synthesis (Fig. 2B). This suggests that CP<sub>CPm</sub> most likely initiates from the upstream CUG and contains additional glutamic acid and threonine residues at its N-terminus (Fig. 3A). The presence of these two additional residues correlates with the slightly slower mobility of CP<sub>CPm</sub> compared with CP<sub>WT</sub> (Figs. 2B, 3B, and 4B). TCV-CPm3, a mutant virus with glutamic acid and threonine codons inserted after the wt CP initiation codon, synthesized wt levels of a CP that likely has the same two residues at its N-terminus as CP<sub>CPm</sub> and comigrates with CP<sub>CPm</sub> on denaturing polyacrylamide gels (Figs. 3A and 3B). Although TCV-CPm3 mimicked TCV-CPm in overcoming the resistance of Di-0 to TCV, symptoms were no longer attenuated by sat-RNA C (Fig. 3C). This suggests that the two additional residues are not sufficient to induce symptom attenuation by sat-RNA C.

An alternative explanation, which cannot be ruled out, is that we have not yet determined the correct initiation codon for CP<sub>CPm</sub> and that alteration of the CUG destabilizes the CP. However, this possibility is unlikely because no known noncanonical initiation codons are found any farther upstream before encountering an in-frame termination codon.

To determine whether the reduced levels of CP found in protoplasts infected with TCV-CPm and CPm-L are responsible for symptom attenuation by sat-RNA C, wt TCV genomic RNA was altered to produce less CP<sub>WT</sub> mRNA (the 1.45-kb sgRNA). TCV-SGm synthesized reduced levels of CP<sub>WT</sub> in protoplasts, and symptoms were partially attenuated by sat-RNA C in both ecotypes of *A. thaliana* (Fig. 4C). This was in slight contrast with TCV-CPm, whose symptoms could be completely attenuated by sat-RNA C in Col-0 and partially attenuated in Di-0 (Figs. 2C, 3C, and 4C). These results suggest that the mutant nature of CP<sub>CPm</sub> contributes to the ability of sat-RNA C to completely attenuate viral symptoms.

The TCV CP is dispensable for replication in proto-

plasts (Hacker *et al.*, 1992; Kong *et al.*, 1995), a result confirmed by the near wt accumulation of CPm-T genomic RNA in protoplasts in the absence of detectable CP synthesis (Fig. 2B). For mutant viruses synthesizing reduced levels of CP (TCV-CPm and CPm-L), the ratios of CP to CP mRNA (the 1.45-kb sgRNA) quantified using densitometry were similar (92% versus 90%). This result suggests that the increased accumulation of CP<sub>CPm-L</sub> compared with CP<sub>CPm</sub> in infected protoplasts was due to an increase in virus replication. Because CPm-L produced 35% of wt levels of CP, the inability to detect virions suggests that as with TCV-CPm, either insufficient CP is present for virion nucleation and/or the presence of mutations in the N-terminal region of the CP inhibits virion formation. The RNA binding activity of TCV CP during encapsidation has been suggested to result from electrostatic interactions between the RNA and the N-terminal basic R domain of the CP, which carries a considerable positive charges (Carrington *et al.*, 1987). CP<sub>CPm</sub> and CP<sub>CPm3</sub> likely have an extra negatively charged residue (E) at the CP N-terminus (Figs. 2A and 3A), which could compromise electrostatic interactions, resulting in less encapsidation-competent CP. This could explain why only 80% of wt levels of virions were present in TCV-CPm3-infected protoplasts despite the wt levels of viral RNA and CP (Fig. 3B). Furthermore, efficient encapsidation requires a highly cooperative assembly interaction between the CP and viral RNA (Skuzeski and Morris, 1995). Our inability to detect more than trace amounts of TCV-SGm virions (Fig. 4B) suggests that the amount of CP<sub>WT</sub> synthesized (20% of wt) is near the threshold required for virion formation. Because comparable levels of mutant CP resulted in no detectable virions, near-threshold amounts of encapsidation-impaired CP present in cells infected with TCV-CPm are likely what restrict the formation of virions to undetectable levels (Figs. 2B, 3B, and 4B).


With only a few exceptions (Gardiner *et al.*, 1988; Petty and Jackson, 1990; Dalmay *et al.*, 1992; Scholthof *et al.*, 1993; Xiong *et al.*, 1993; Azzam *et al.*, 1994), viral CPs are required for systemic infection of plant viruses (Dawson *et al.*, 1988; Sacher and Ahlquist, 1989; Allison *et al.*, 1990; Chapman *et al.*, 1992; Forster *et al.*, 1992; Hacker *et al.*, 1992; Sit and AbouHaidar, 1993; Dolja *et al.*, 1994; Bransom *et al.*, 1995). However, assembly of CP units into virions may not be necessary for movement of all viruses. Although some studies indicate that virion formation is essential for virus long-distance movement (Saito *et al.*, 1990; Schmitz and Rao, 1998), other reports (Dalmay *et al.*, 1992; Dolja *et al.*, 1994; Schneider *et al.*, 1997; Kaplan *et al.*, 1998), including this study, indicate that although the CP is required, virions appear to be dispensable.

Recent studies (Kong *et al.*, 1997a) indicate that the 3'-terminal 100 bases of sat-RNA C, which is 90% similar to the analogous region in TCV genomic RNA, is the

sat-RNA region involved in symptom modulation. The disappearance or amelioration of TCV symptoms in the presence of sat-RNA C may therefore seem to echo the cosuppression phenomenon found in transgenic plants. Cosuppression occurs when the introduction of a transgene encoding part of or the entire coding sequence of a particular host gene can suppress expression of the transgene and the endogenous host gene (reviewed by Montgomery and Fire, 1998). However, the mechanism underlying symptom attenuation by sat-RNA C is unlikely to be due to cosuppression for the following reasons: first, sat-RNA C-mediated resistance of *A. thaliana* to TCV-CPm only moderately affects the level of genomic RNA in inoculated leaves (Kong *et al.*, 1997b), which is different from the substantial amount of RNA degradation during cosuppression (Lee *et al.*, 1997); second, another normally virulent subviral RNA, defective interfering (DI) RNA G, has greater sequence similarity (94%) with the TCV genomic RNA sequence in the 3'-terminal region (Kong *et al.*, 1997a). However, although DI RNA G attenuates the symptoms of TCV-CP<sub>CCFV</sub>, it does not attenuate the symptoms of TCV-CPm.

Different sat-RNAs may use different mechanisms to attenuate symptoms. In many hosts, disease attenuation by CMV sat-RNAs is accompanied by a reduction in virus accumulation (Kaper and Tousignant, 1977; Kaper and Collmer, 1988). In contrast, CMV sat-RNA symptom attenuation of tomato aspermy virus is not always accompanied by a noticeable decrease in the level of viral RNAs (Moriones *et al.*, 1992). Although the attenuation of TCV-CP<sub>CCFV</sub> symptoms by sat-RNA C was associated with a substantial reduction in virus replication (Kong *et al.*, 1997b), sat-RNA C attenuation of TCV-CPm symptoms did not involve a large reduction in virus accumulation but rather was associated with a reduction in virus movement (Kong *et al.*, 1997b). Because virions are not detected in TCV-CPm-infected protoplasts, some other type of viral RNA-CP complex could be engaged in virus movement. Symptom attenuation by sat-RNA C of TCV variants producing reduced levels of CP could therefore result from sat-RNA C competing with the viral genomic RNA for limited amounts of CP, thus reducing or eliminating genomic RNA-CP complexes required for systemic movement of the virus.

A second possibility for how sat-RNA C may attenuate symptoms involves sequestration of factor or factors required for virus movement. Recently, studies have led to suggestions that host factors are involved in trafficking viruses from phloem parenchyma cells into phloem sieve elements and back out, steps required for long-distance virus movement (Schaad and Carrington, 1996; Gilbertson and Lucas, 1996). The 3' end region of sat-RNA C that has recently been shown to be responsible for symptom attenuation (Wang and Simon, manuscript in preparation) may be targeted by either the viral CP or a putative host factor (X) that is involved in virus long-



**FIG. 5.** Putative model for symptom modulation by sat-RNA C in TCV/*Arabidopsis*. In this model, the TCV CP (indicated by small gray circles) binds directly (or indirectly) to the 3' end of sat-RNA C (satC) but not sat-RNA D (satD) and competes for binding with a putative host factor "X" (indicated by large black circles) that is involved in virus long-distance movement. (A) When high levels of CP are present, CP outcompetes "X" for binding to the 3' end of satC, leaving "X" available for virus long-distance movement (i.e., systemic infection). (B) When low levels of CP are present, "X" outcompetes the CP for binding to the 3' end of satC, and sequestration of "X" by satC restricts virus movement and results in attenuation of symptoms. (C) "X" outcompetes non-TCV CP (indicated by small hatched circles), as present in TCV-CP<sub>CCFV</sub>-infected plants, for binding to the 3' end of satC, resulting in similar symptom modulation as shown in B.

distance movement (Fig. 5). Regardless of the nature of the interaction between the CP and the 3' region of sat-RNA C (i.e., direct or indirect), the presence of high levels of viral CP (as in TCV- or TCV-CPm3-infected plants) may exclude the binding of X to the 3' end of the sat-RNA C. Therefore, the putative X is available to help viral RNA move systemically through the plant (Fig. 5A). In contrast, in the presence of reduced levels of CP (as in TCV-CPm-, CPm-L-, or TCV-SGm-infected plants), X is able to outcompete the CP for binding to the 3'-terminus of sat-RNA C, and sequestration of X by sat-RNA C could lead to restriction of virus long-distance movement and result in symptom attenuation (Fig. 5B). On the other hand, when TCV-CP<sub>CCFV</sub> is the helper virus, the CCFV CP (which shares only 65% identity with CP<sub>WT</sub>; Oh *et al.*, 1995) may not recognize the 3' end of sat-RNA C, allowing X to bind and resistance to be mediated (Fig. 5C). Studies to distinguish between possible models are currently under way.

The susceptibility of Di-0 to TCV-CPm and TCV-CPm3 suggests that the N-terminus of the CP is important for the resistance of Di-0 to TCV. In addition, the susceptibility of Di-0 to TCV-SGm suggests that reducing the levels of wt CP can also eliminate resistance. In addition to its role in resistance, the TCV CP also contributes to expression of symptoms. The symptoms of TCV-CPm3

TABLE 1  
Summary of the Oligonucleotides Used in the Study

Application/construct	Name	Position in TCVms	Sequence <sup>a</sup>	Polarity <sup>b</sup>
Site-directed mutagenesis	CPm-LTO	2737–2756	5'- <u>AYTGGAAA</u> VGGAAAATGATC-3'	+
	CPm3C(-)	2737–2762	5'-ACTGGAAATGG <u>AAACT</u> GAAAATGATC-3'	+
	OL3270C(+)	3270–3287	5'-TCCAGGGCACGCTAGATA-3'	-
	OL2736C(+)	2717–2736	5'-GTTGATGCTTATGTGTTGCT-3'	-
	1.7 LB	2243–2260	5'-ACCGGGGGTTCGGCTACA-3'	+
	SGmC(+)	2584–2603	5'-GGGGA <u>T</u> GTCGGGCACTGACA-3'	-
	OL2604C(-)	2604–2623	5'-GTGGGTAAATATATGCTTTCT-3'	+
	RV-1C(-)	2095–2114	5'- <u>GATAT</u> CTTGCCTGAAGAGGAATTT-3'	+
RNA gel blots	OL3892C(+)	3893–3913	5'-CCGTTTTTGGTCCCTAACACA-3'	-

<sup>a</sup> Underlined sequences are inserted or mutated sequences.

<sup>b</sup> Polarity refers to homology (+) or complementarity (-) with plus strands of TCV genomic RNA.

were more severe than those of TCV in Col-0, even though similar levels of viral RNA and CP were synthesized in protoplasts (Fig. 3). This suggests that mutations at the N-terminus of the CP exacerbate symptoms. The multifunctional nature of viral CPs may be due to multiple determinants that can be hidden (or exposed) by changes in the structure or quantity of the protein. This property of CPs has been demonstrated for tobacco mosaic virus: a reduction of subunit interactions was hypothesized to expose a site on the CP that interacts with the N' gene product of tobacco, resulting in hypersensitive resistance (Culver *et al.*, 1994). It is also possible that virions themselves elicit more severe symptoms, whereas the CP monomer does not elicit such a response. This could explain why TCV, TCV-CPm3 (Fig. 3), and TCV-SGm (Fig. 4) caused more severe symptoms than TCV-CPm and CPm-L (Fig. 2).

## MATERIALS AND METHODS

### Virus strains and plasmid constructions

Plasmids containing full-length cDNAs of TCV (pT7TCVms; Oh *et al.*, 1995), sat-RNA C [pT7satC(+); Song and Simon, 1994], and TCV-CPm (pT7TCV-CPm; Kong *et al.*, 1997) downstream from a T7 RNA polymerase promoter have been described.

For construction of plasmids CPm-L, CPm-T, and CPm-O, oligonucleotide CPm-LTO, containing degenerate nucleotides in its sequence (all the oligonucleotides are listed in Table 1), was used with oligonucleotide OL3270C(+) in a polymerase chain reaction (PCR) using pT7TCV-CPm as template. A second PCR was performed with oligonucleotides 1.7LB and OL2736C(+). The PCR products were digested with *BsmI* and *EcoRI*, respectively, and the larger fragments were purified and combined. The mixture was then ligated to pT7TCVms that had been previously digested with *BsmI* and *EcoRI*. Plasmid TCV-CPm3 was generated in a similar fashion except that oligonucleotide CPm-LTO was replaced with CPm3C(-). Plas-

mid TCV-SGm was generated in a similar fashion except that the template and oligonucleotide CPm-LTO were replaced with pT7TCVms and OL2604C(-), respectively, and the second PCR was performed with oligonucleotides SGmC(+) and RV-1C(-).

### Plant growth and inoculations

Plants (*A. thaliana* ecotypes Col-0 and Di-0) were grown in growth chambers at 20°C as described by Li and Simon (1990). Plant seedlings at the six- to eight-leaf stage were mechanically inoculated on the oldest leaf pair as described previously (Kong *et al.*, 1997b) with 0.1 mg/ml full-length transcripts synthesized *in vitro* from cloned cDNAs using T7 RNA polymerase (Carpenter *et al.*, 1995). For experiments examining the effects of sat-RNA C on symptom modulation, 0.01 mg/ml of full-length transcripts synthesized from the cloned cDNA of sat-RNA C was included in the inoculum.

### Preparation and inoculation of *A. thaliana* protoplasts

Protoplasts were prepared from Col-0 callus cultures as described (Kong *et al.*, 1997b). Protoplasts ( $5 \times 10^6$ ) were inoculated with 20 µg of genomic RNA transcripts synthesized *in vitro* as previously described (Kong *et al.*, 1997b).

### Protein gel blot analysis

Total proteins were extracted from protoplasts by vortexing the cells in an equal volume of extraction buffer [125 mM Tris-HCl, pH 6.8, 0.1% SDS, and 20% glycerol (v/v)] followed by centrifugation at 10,000 rpm for 5 min in a microcentrifuge to collect the supernatants. Total protein or isolated virions were separated on either 12% SDS-polyacrylamide gels or 1% agarose gels, respectively, containing 50 mM Tris base/38 mM glycine, pH 8.3, as previously described (Heaton, 1992). Total protein or virions were transferred to NitroPlus membrane (Micron Separations Inc., Westborough, MA) as previously de-

scribed (Kong *et al.*, 1997b). Protein gel blot analysis was performed as described by Ausubel *et al.* (1987) with some modifications as described by Kong *et al.* (1997b).

### Virion isolation and analysis

Virus particles were isolated from infected protoplasts as previously described (Qu and Morris, 1997) with modifications described by Kong *et al.* (1997b). Virus particles were analyzed by electrophoresis through 1% agarose gels prepared in 50 mM Tris base/38 mM glycine, pH 8.3, as previously described (Laakso and Heaton, 1993) followed by protein gel blot analysis as described above.

### RNA gel blot analysis

Four micrograms of total RNA isolated from protoplasts (Simon *et al.*, 1992) were denatured by heating in 50–70% formamide and then subjected to electrophoresis through nondenaturing 1.5% agarose gels. RNA was then transferred to NitroPlus membrane (Micron Separations Inc.) and subjected to hybridization with an oligonucleotide probe specific for TCV (Table 1) or a probe specific for plant ribosomal RNAs (Simon *et al.*, 1992) as previously described (Wang and Simon, 1997).

### ACKNOWLEDGMENT

This work was supported by National Science Foundation Grants MCB-9630191 and MCB-9728277 (A.E.S.).

### REFERENCES

- Allison, R., Thompson, C., and Ahlquist, P. (1990). Regeneration of a functional RNA virus genome by recombination between deletion mutants and requirement for cowpea chlorotic mottle virus 3a and coat genes for systemic infection. *Proc. Natl. Acad. Sci. USA* **87**, 1820–1824.
- Ausubel, F. M., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A., and Struhl, K., Eds. (1987). "Current Protocols in Molecular Biology." New York, John Wiley and Sons.
- Azzam, O., Frazer, J., de La Rosa, D., Beaver, J. S., Ahlquist, P., and Maxwell, D. P. (1994). Whitefly transmission and efficient ssDNA accumulation of bean golden mosaic geminivirus require functional coat protein. *Virology* **204**, 289–296.
- Baulcombe, D., Devic, M., Jaegle, M., and Harrison, B. (1988). Control of viral infection in transgenic plants by expression of satellite RNA of cucumber mosaic virus. In "UCLA Symposium on Molecular and Cellular Biology," New Series, Vol. 101 (Staskowicz, B., Ahlquist, P., and Yoder, O., Eds.), Alan R. Liss, Inc., New York.
- Beames, B., Braunagel, S., Summers, M. D., and Lanford, R. E. (1991). Polyhedrin initiator codon altered to AUU yields unexpected fusion protein from a baculovirus vector. *BioTechniques* **11**, 378–383.
- Blok, V. C., Ziegler, A., Robinson, D. J., and Murrant, A. F. (1994). Sequences of 10 variants of the satellite-like RNA3 of groundnut rosette virus. *Virology* **202**, 25–32.
- Bransom, K. L., Weiland, J. J., Tsai, C.-H., and Dreher, T. W. (1995). Coding density of the turnip yellow mosaic virus genome: Roles of the overlapping coat protein and p206-readthrough coding regions. *Virology* **206**, 403–412.
- Carpenter, C. D., Oh, J.-W., Zhang, C., and Simon, A. E. (1995). Involvement of a stem-loop structure in the location of junction sites in viral RNA recombination. *J. Mol. Biol.* **245**, 608–622.
- Carrington, J. C., Heaton, L. A., Zuidema, D., Hillman, B. I., and Morris, T. J. (1989). The genome structure of turnip crinkle virus. *Virology* **170**, 219–226.
- Carrington, J. C., Morris, T. J., Stockley, P. G., and Harrison, S. C. (1987). Structure and assembly of turnip crinkle virus. IV. Analysis of the coat protein gene and implications of the subunit primary structure. *J. Mol. Biol.* **194**, 265–276.
- Celix, A., Rodriguez-Cerezo, E., and Garcia-Arenal, F. (1997). New satellite RNAs, but no DI RNAs, are found in natural population of tomato bushy stunt tomosvirus. *Virology* **239**, 277–284.
- Chapman, S., Kavanagh, T., and Baulcombe, D. C. (1992). Potato virus X as a vector for gene expression in plants. *Plant J.* **2**, 549–557.
- Collmer, C. W., and Howell, S. H. (1992). Role of satellite RNA in the expression of symptoms caused by plant viruses. *Annu. Rev. Phytopathol.* **30**, 419–442.
- Culver, J. N., Stubbs, G., and Dawson, W. (1994). Structure-function relationship between tobacco mosaic virus coat protein and hypersensitivity in *Nicotiana sylvestris*. *J. Mol. Biol.* **242**, 130–138.
- Curran, J. A., and Kolakofsky, D. (1988). Ribosomal initiation from an ACG codon in the Sendai virus P/C mRNA. *EMBO J.* **7**, 245–251.
- Dalmay, T., Rubino, L., Burgyan, J., and Russo, M. (1992). Replication and movement of a coat protein mutant of cymbidium ringspot tomosvirus. *Mol. Plant-Microbe Interact.* **5**, 379–383.
- Dawson, W. O., Bublick, P., and Grantham, G. L. (1988). Modification of the tobacco mosaic virus coat protein gene affecting replication, movement, and symptomatology. *Phytopathology* **78**, 783–789.
- Dolja, V. V., Haldeman, R., Robertson, N. L., Dougherty, W. G., and Carrington, J. C. (1994). Distinct function of capsid protein in assembly and movement of tobacco etch potyvirus. *EMBO J.* **13**, 1482–1491.
- Forster, R. L. S., Beck, D. L., Guilford, P. J., Voot, D. M., Van Dolleweerd, C. J., and Andersen, M. T. (1992). The coat protein of white clover mosaic potyvirus has a role in facilitating cell-to-cell transport in plants. *Virology* **191**, 480–484.
- Gallie, D. R. (1993). Posttranscriptional regulation of gene expression in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **44**, 77–105.
- Gardiner, W. E., Sunter, G., Brand, L., Elmer, J. S., Rogers, S. G., and Bisaro, D. M. (1988). Genetic analysis of tomato golden mosaic virus: The coat protein is not required for systemic spread or symptom development. *EMBO J.* **7**, 899–904.
- Gerlach, W. L., Llewellyn, D., Haseloff, J. (1987). Construction of a plant disease resistance gene from the satellite RNA of tobacco ringspot virus. *Nature* **328**, 802–805.
- Gilbertson, R. L., and Lucas, W. J. (1996). How do viruses traffic on the 'vascular highway'? *Trends Plant Sci.* **1**, 260–267.
- Gordan, K., Fütterer, J., and Hohn, T. (1992). Efficient initiation of translation at non-AUG triplets in plant cells. *Plant J.* **2**, 809–813.
- Gupta, K. C., and Patwardhan, S. (1988). ACG, the initiator codon for a Sendai virus protein. *J. Biol. Chem.* **263**, 8553–8556.
- Hacker, D. L., Petty, I. T. D., Wei, N., and Morris, T. J. (1992). Turnip crinkle virus genes required for RNA replication and virus movement. *Virology* **186**, 1–8.
- Hann, S. R., King, M. W., Bentley, D. L., Anderson, C. W., and Eisenman, R. N. (1988). A non-AUG translational initiation in *c-myc* exon 1 generates an N-terminally distinct protein whose synthesis is disrupted in Burkitt's lymphomas. *Cell* **52**, 185–195.
- Harrison, B. D., Mayo, M. A., and Baulcombe, D. C. (1987). Virus resistance in transgenic plants that express cucumber mosaic virus satellite RNA. *Nature* **328**, 799–802.
- Heaton, L. A. (1992). Use of agarose gel electrophoresis to monitor conformational changes of some small, spherical plant viruses. *Phytopathology* **82**, 803–807.
- Hogle, J. M., Maeda, K. A., and Harrison, S. C. (1986). Structure and assembly of turnip crinkle virus. I. X-ray crystallographic structure analysis at 3.2 Å resolution. *J. Mol. Biol.* **191**, 625–638.
- Jaegle, M., Devic, M., Longstaff, M., and Baulcombe, D. (1990). Cucumber mosaic virus satellite RNA (Y strain): Analysis of sequence which

- affect yellow mosaic symptoms on tobacco. *J. Gen. Virol.* **71**, 1905–1912.
- Kaper, J. M., and Collmer, C. W. (1988). Modulation of viral plant diseases by secondary RNA agents. In "RNA Genetics: Vol. 3. Variability of RNA Genomes" (Domingo, E., Holland, J. J., and Ahlquist, P., Eds.), pp. 171–194. CRC Press, Boca Raton.
- Kaper, J. M., and Tousignant, M. E. (1977). Cucumber mosaic virus-associated RNA 5. I. Role of host plant and helper strain in determining amount of associated RNA 5 with virions. *Virology* **80**, 186–195.
- Kaper, J. M., Tousignant, M. E., and Geletka, L. M. (1990). Cucumber mosaic virus-associated RNA 5. XII. Symptom modulating effect is codetermined by satellite replication support function of helper virus. *Res. Virol.* **141**, 487–503.
- Kaplan, I. B., Zhang, L., and Palukaitis, P. (1998). Characterization of cucumber mosaic virus. V. Cell-to-cell movement requires capsid protein but not virions. *Virology* **246**, 221–223.
- Kong, Q. (1996). Plant resistance and symptom modulation by a satellite RNA in turnip crinkle virus/*Arabidopsis* system. PhD dissertation, University of Massachusetts at Amherst.
- Kong, Q., Oh, J.-W., Carpenter, C. D., and Simon, A. E. (1997a). The coat protein of turnip crinkle virus is involved in subviral RNA-mediated symptom modulation and accumulation. *Virology* **238**, 478–485.
- Kong, Q., Oh, J.-W., and Simon, A. E. (1995). Symptom attenuation by a normally virulent satellite RNA of turnip crinkle virus is associated with the coat protein open reading frame. *Plant Cell* **7**, 1625–1634.
- Kong, Q., Wang, J., and Simon, A. E. (1997b). Satellite RNA-mediated resistance to turnip crinkle virus in *Arabidopsis* involves a reduction in virus movement. *Plant Cell* **9**, 2051–2063.
- Kurath, G., and Palukaitis, P. (1989). Satellite RNAs of cucumber mosaic virus: Recombinants constructed *in vitro* reveal independent functional domains for chlorosis and necrosis in tomato. *Mol. Plant-Microbe Interact.* **2**, 91–96.
- Laakso, M. M., and Heaton, L. A. (1993). Asp → Asn substitutions in the putative calcium-binding site of the turnip crinkle virus coat protein affect virus movement in plants. *Virology* **197**, 774–777.
- Lee, K. Y., Baden, C., Howie, W. J., Bedbrook, J., and Dunsmuir, P. (1997). Post-transcriptional gene silencing of ACC synthase in tomato results from cytoplasmic RNA degradation. *Plant J.* **12**, 1127–1137.
- Li, W.-Z., Qu, F., and Morris, T. J. (1998). Cell-to-cell movement of turnip crinkle virus is controlled by two small open reading frames that function *in trans*. *Virology* **244**, 405–416.
- Li, X. H., and Simon, A. E. (1990). Symptom intensification on cruciferous hosts by the virulent satellite RNA of turnip crinkle virus. *Phytopathology* **80**, 238–242.
- Masuta, C., and Takanami, Y. (1989). Determination of sequence and structural requirements for pathogenicity of a cucumber mosaic virus satellite RNA (Y-satRNA). *Plant Cell* **1**, 1165–1173.
- Mehdi, H., Ono, E., and Gupta, K. C. (1990). Initiation of translation at CUG, GUG, and ACG codons in mammalian cells. *Gene* **91**, 173–178.
- Militao, V., Moreno, I., Rodriguez-Cerezo, E., and Garcia-Arenal, F. (1998). Differential interactions among isolates of peanut stunt cucumovirus and its satellite RNA. *J. Gen. Virol.* **79**, 177–184.
- Montgomery, M. K., and Fire, A. (1998). Double-stranded RNA as a mediator in sequence-specific genetic silencing and co-suppression. *Trends Genet.* **14**, 255–258.
- Moriones, E., Diaz, I., Rodriguez-Cerezo, E., Fraile, A., and Garcia-Arenal, F. (1992). Differential interactions among strains of tomato aspermy virus and satellite RNAs of cucumber mosaic virus. *Virology* **186**, 475–480.
- Murant, A. F., and Kumar, I. K. (1990). Different variants of the satellite RNA of groundnut rosette virus are responsible for the chlorotic and green forms of groundnut rosette disease. *Ann. Appl. Biol.* **117**, 85–92.
- Naidu, R. A., Collins, G. B., and Ghabrial, S. A. (1992). Peanut stunt virus satellite RNA: Analysis of sequences that affect symptom attenuation in tobacco. *Virology* **189**, 668–677.
- Oh, J.-W., Kong, Q., Song, C., Carpenter, C. D., and Simon, A. E. (1995). Open reading frame of turnip crinkle virus involved in satellite symptom expression and incompatibility with *Arabidopsis thaliana* ecotype Dijon. *Mol. Plant-Microbe Interact.* **8**, 979–987.
- Oncino, C., Hemmer, O., and Fritsch, C. (1995). Specificity in the association of tomato black ring virus satellite RNA with helper virus. *Virology* **213**, 87–96.
- Palukaitis, P. (1988). Pathogenicity regulation by satellite RNAs of cucumber mosaic virus: Minor nucleotide sequence changes alter host response. *Mol. Plant-Microbe Interact.* **1**, 175–181.
- Peabody, D. S. (1989). Translation initiation at non-AUG triplets in mammalian cells. *J. Biol. Chem.* **264**, 5031–5035.
- Petty, I. T. D., and Jackson, A. O. (1990). Mutational analysis of barley stripe mosaic virus RNA  $\beta$ . *Virology* **179**, 712–718.
- Reynolds, J. E., Kaminski, A., Kettinen, H. J., Grace, K., Clarke, B. E., Carroll, A. R., Rowlands, D. J., and Jackson, R. J. (1995). Unique features of internal initiation of hepatitis C virus RNA translation. *EMBO J.* **14**, 6010–6020.
- Rodriguez-Alvarado, G., and Roossinck, M. J. (1997). Structural analysis of a necrogenic strain of cucumber mosaic cucumovirus satellite RNA in planta. *Virology* **236**, 155–166.
- Rohde, W., Gramstat, A., Schmitz, J., Tacke, E., and Prüfer, D. (1994). Plant viruses as model systems for the study of non-canonical translation mechanisms in higher plants. *J. Gen. Virol.* **75**, 2141–2149.
- Roossinck, M. J., Sleat, D., and Palukaitis, P. (1992). Satellite RNAs of plant viruses: Structures and biological effects. *Microbiol. Rev.* **56**, 265–279.
- Sacher, R., and Ahlquist, P. (1989). Effects of deletions in the N-terminal basic arm of brome mosaic virus coat protein on RNA packaging and systemic infection. *J. Virol.* **63**, 4545–4552.
- Saito, T., Yamanaka, K., and Okada, Y. (1990). Long-distance movement and viral assembly of tobacco mosaic virus mutants. *Virology* **176**, 329–336.
- Saris, C. J. M., Domen, J., and Berns, A. (1991). The *pim-1* oncogene encodes two unrelated protein-serine/threonine kinases by alternative initiation at AUG and CUG. *EMBO J.* **10**, 655–664.
- Schaad, M. C., and Carrington, J. C. (1996). Suppression of long-distance movement of tobacco etch virus in a nonsusceptible host. *J. Virol.* **70**, 2556–2561.
- Schmitz, I., and Rao, A. L. N. (1998). Deletions in the conserved amino-terminal basic arm of cucumber mosaic virus coat protein disrupt virion assembly but do not abolish infectivity and cell-to-cell movement. *Virology* **248**, 323–331.
- Schmitz, J., Prüfer, D., Rohde, W., and Tacke, E. (1996). Non-canonical translation mechanisms in plants: Efficient *in vitro* and in planta initiation at AUU codons of the tobacco mosaic virus enhancer sequence. *Nucleic Acids Res.* **24**, 257–264.
- Schneider, W. L., Greene, A. E., and Allison, R. F. (1997). The carboxy-terminal two-thirds of the cowpea chlorotic mottle bromovirus capsid is incapable of virion formation yet supports systemic movement. *J. Virol.* **71**, 4862–4865.
- Scholthof, H. B., Morris, T. J., and Jackson, A. O. (1993). The capsid protein of tomato bushy stunt virus is dispensable for systemic movement and can be replaced for localized expression of foreign genes. *Mol. Plant-Microbe Interact.* **6**, 309–322.
- Simon, A. E. (1988). Satellite RNAs of plant viruses. *Plant Mol. Biol. Reporter* **6**, 240–252.
- Simon, A. E., Engel, H., Johnson, R. P., and Howell, S. H. (1988). Identification of regions affecting virulence, RNA processing and infectivity in the virulent satellite of turnip crinkle virus. *EMBO J.* **7**, 2645–2651.
- Simon, A. E., Li, X. H., Lew, J. E., Stange, R., Zhang, C., Polacco, M., and Carpenter, C. D. (1992). Susceptibility and resistance of *Arabidopsis thaliana* to turnip crinkle virus. *Mol. Plant-Microbe Interact.* **5**, 496–50.
- Sit, T. L., and AbouHaidar, M. G. (1993). Infectious RNA transcripts derived from cloned cDNA of papaya mosaic virus: Effect of muta-

- tions to the capsid and polymerase proteins. *J. Gen. Virol.* **74**, 1133–1140.
- Skuzeski, J. M., and Morris, T. J. (1995). Quantitative analysis of the binding of turnip crinkle virus coat protein to RNA fails to demonstrate binding specificity but reveals a highly cooperative assembly interaction. *Virology* **210**, 82–90.
- Sleat, D. E., and Palukaitis, P. (1990a). Site-directed mutagenesis of a plant viral satellite RNA changes its phenotype from ameliorative to necrogenic. *Proc. Natl. Acad. Sci. USA* **87**, 2946–2950.
- Sleat, D. E., and Palukaitis, P. (1990b). Induction of tobacco chlorosis by certain cucumber mosaic virus satellite RNAs is specific to subgroup II helper strain. *Virology* **176**, 292–295.
- Sleat, D. E., Zhang, L., and Palukaitis, P. (1994). Mapping determinants within cucumber mosaic virus and its satellite RNA for the induction of necrosis in tomato plants. *Mol. Plant-Microbe Interact.* **7**, 189–195.
- Song, C., and Simon, A. E. (1994). RNA-dependent RNA polymerase from plants infected with turnip crinkle virus can transcribe (+)- and (–)-strands of virus-associated RNAs. *Proc. Natl. Acad. Sci. USA* **91**, 8792–8796.
- Takanami, Y. (1981). A striking change in symptoms on cucumber mosaic virus-infected tobacco plants induced by a satellite RNA. *Virology* **109**, 120–126.
- Taliansky, M. E., and Robinson, D. J. (1997). Trans-acting untranslated elements of groundnut rosette virus satellite RNA are involved in symptom production. *J. Gen. Virol.* **78**, 1277–1285.
- Tien, P., and Wu, G. (1991). Satellite RNA for the biocontrol of plant disease. *Adv. Virus Res.* **39**, 321–329.
- Wang, J., Carpenter, C. D., and Simon, A. E. (1999). Minimal sequence and structural requirements for a subgenomic RNA promoter for turnip crinkle virus. *Virology* **253**, 327–336.
- Wang, J., and Simon, A. E. (1997). Analysis of the two subgenomic RNA promoters for turnip crinkle virus *in vivo* and *in vitro*. *Virology* **232**, 174–186.
- Xiong, Z., Kim, K. H., Giesman-Cookmeyer, D., and Lommel, S. A. (1993). The roles of the red clover necrotic mosaic virus capsid and cell-to-cell movement proteins in systemic infection. *Virology* **192**, 27–32.
- Zhang, L., Kim, C. H., and Palukaitis, P. (1994). The chlorosis-induction domain of the satellite RNA of cucumber mosaic virus: Identifying sequences that affect accumulation and the degree of chlorosis. *Mol. Plant-Microbe Interact.* **7**, 208–213.